

Cosmology with variable G and nonlinear electrodynamics

G W Joseph and A Övgün* 

Physics Department, Eastern Mediterranean University, via Mersin 10, 99628 Famagusta, North Cyprus, Turkey

Received: 10 October 2020 / Accepted: 05 April 2021

Abstract: In a bid to resolve lingering problems in cosmology, more focus is being tilted towards cosmological models in which physical constants of nature are not necessarily real constants but vary with cosmic time. In this paper, we study a cosmological model in nonlinear electrodynamics with Newton's gravitational constant G , which is not a constant but varies in terms of a power law of the scale factor of the universe. Hence, the evolution of the scale factor $a(t)$ is studied in this model, which depends on a fine-tuning term of nonlinear electrodynamics, α . Finally, we verify the stability of the model using the speed of sound.

Keywords: Cosmology; Nonlinear electrodynamics; Inflation; Acceleration of the universe; Causality; Classical stability; Variable G

1. Introduction

Remarkable achievements recorded in the field of cosmology in the past decades can be attributed to observations of cosmic microwave background (CMB) radiation and type-Ia supernovae, suggesting that cosmic expansion is accelerating [1]. It is well-known that Maxwell's electrodynamics, as a source of the Friedmann-Robertson-Walker spacetime, which is the central to the standard cosmological model (SCM), causes an initial singularity: the breakdown in geometrical structure of space and time at a finite time in the past. This initial singularity around the very early universe has revealed an unpredictable large value of the energy density and curvature of spacetime, thus departing from the domain of the SCM. This setback has given rise to other secondary cosmological problems, such as the horizon, flatness, and monopole problems. The secondary problems have been resolved by using scalar fields. Moreover, magnetic universe models, which portray no initial singularity due to strong electromagnetic field in the modified nonlinear electrodynamics (NED) in the early universe, have been used in the literature to resolve this puzzle [2–18]. The introduction of cosmological constant Λ , or scalar field with a type of potential function or modifying the theory of gravity with the aid of some

function $f(r)$, can effectively explain the acceleration of the universe [19].

All these models are not without their own setbacks [20–23]. For instance, the smallness of the cosmological constant when compared with the vacuum energy is very difficult to explain. Furthermore, the choice of $f(r)$ function in the modified gravity is not unique. Moreover, inflationary cosmological models provide convincing solutions to the horizon problem, flatness problem, and small value of cosmological constant, no model of inflation has been experimentally confirmed.

In scalar-tensor theory, much potential leads to cosmic inflation and acceleration of the universe. When NED is coupled to a gravitational field, it can give the necessary negative pressure and enhance cosmic inflation. Therefore, it is interesting to explore a new model of NED with variable gravitational constant where stochastic magnetic field drives the inflation of the universe.

Magnetic fields at different levels in the past have effects on various cosmological and astrophysical phenomena observed in the universe. Throughout the universe, physical processes over a large range of scales are influenced by magnetic fields in the universe. The present and associate magnetic fields could therefore arise from the flow of materials in the high-temperature vicinity of the big bang. This corresponds to the primordial magnetic fields (PMF). Adequate knowledge of the PMF offers information about the early universe between the inflationary and recombination eras. Such magnetic fields originate from

*Corresponding author, E-mail: ali.ovgun@emu.edu.tr

cosmic phase transition with the bluer spectrum and vector potential produced during the inflation era. Stochastic inconsistencies of EM field in relativistic electron-positron plasma produce PMF from thermal instabilities in the pre-recombination plasma. The plasma sustains the magnetic fluctuations in the early universe before the era of Big Bang nucleosynthesis during the radiation-dominated era.

Using the new type of NED containing a stochastic magnetic background with nonvanishing B^2 gives cosmological models void of initial singularities. Other cosmological models lacking a primordial singularity, such as non-minimal coupling, quantum gravity effect, Lagrangian with quadratic term, inflation by scalar fields, and NED without modification of general relativity, have been introduced in the literature to solve the puzzle of cosmology and mystery of inflation [19, 24–44].

Max Born and Leopold Infeld used the idea of Gustav who in 1909 began constructing a pure electromagnetic theory of charged particles to propose a new theory in 1934: fully relativistic and gauge-invariant nonlinear electrodynamics [45]. Born-Infeld proposed a nonlinear fields Lagrangian with an interesting attribute of transforming to Maxwell's theory for low electromagnetic fields. Since there are no new degrees of freedoms, such as scalar fields or branes, works of cosmology described by NED should have some interesting features of cosmic importance. The sources of cosmic inflation can be traced to nonlinear electromagnetic radiation which is explained by modified Maxwell's equations. When coupled with the gravitational fields, NED may give negative pressure and also can lead to cosmic inflation [19]. The evolution of the universe, when explored with a new NED model such that electromagnetic field coupled to gravitational field prevents cosmic singularity at the big bang. The electromagnetic and gravitational fields were very strong during the evolution of the early universe, thereby leading to quantum correction and giving birth to NED [27–29]. One can present a model of the universe filled with a Born-Infeld type of nonlinear electromagnetic field, which is inhomogeneous and anisotropic without singularity.

Recently, interest is being geared towards cosmological models in which physical constants of nature are varying with time [46–48]. For instance, in the varying speed of light (VSL) theories, where the speed of light is spatial, temporal and wavelength-dependent, most pending problems of standard cosmological models are being resolved without considering inflation [49–51]. The VSL theories, which deal with the shortcomings of inflation, have not address issues related to the initial big bang singularity. The Newtonian gravitational constant G occurs in the source term of Einstein's field equation of the general theory of relativity, which is a fundamental equation for

developing every model of cosmology. In Einstein's field equation, G acts as a coupling constant between the geometry of spacetime and matter. In quantum mechanics, G is essential in the definition of the Planck constant [52]. While in SCM, G is an invariant quantity. It has been noted that there is significant evidence that the gravitational constant G can vary in time [53]. Dirac, in 1937, argued that variations in G of about 5 parts in 10^{11} per annum could explain the relatively small strength of the gravitational force compared to other fundamental forces of nature. In order to unify gravitation and elementary particle physics, Einstein's theory with time-varying G is already in the literature [50, 54–65]. Singh [57] has presented a cosmological model with $G \sim a^{2/\alpha}$, where a is the scale factor of the universe and α a constant. When either G or the cosmological constant Λ is varied with time, the Einstein field equation is still preserved. It has been shown that, the variable Newtonian gravitation constant can account for dark energy and most of its effects, and current dynamical dark energy models using time-dependent cosmological constant terms are being considered [58].

In this paper, we study a model involving nonlinear electrodynamics coupled to general relativity, additionally assuming a time-dependent gravitational constant, and study its cosmological dynamics, in order to unify the different epochs during the evolution of the universe. We show that variable G in the study of nonlinear electromagnetic radiation is a source of inflation in the early universe.

The structure of the paper is as follows: in Sect. 2, we briefly introduce the cosmology of a universe filled with nonlinear magnetic monopole fields. In Sect. 3, we obtain the evolution of the universe filled with nonlinear magnetic monopole fields and variable gravitational constant. In Sect. 4, we check the stability of the model and give our conclusion in Sect. 5.

2. Nonlinear magnetic monopole fields and cosmology

In nonlinear electrodynamics, we define the Lagrangian density by [10]

$$\mathcal{L}_{\text{NED}} = -\frac{\mathcal{F}^\alpha}{4}, \quad (1)$$

where α is the electromagnetic fine structure constant and \mathcal{F} denotes an invariant quantity known as the Maxwell invariant. Since the matter part of the Lagrangian is independent of the metric's derivatives, in tensorial language the matter energy-momentum definition using (1) is given as [12]

$$T_{\mu\nu} = -K_{\mu\lambda} F_{\nu}^{\lambda} + g_{\mu\nu} \mathcal{L}_{\text{NED}}, \quad (2)$$

with

$$K_{\mu\lambda} = \frac{\partial \mathcal{L}_{\text{NED}}}{\partial \mathcal{F}} F_{\mu\lambda}, \quad (3)$$

where $g_{\mu\nu}$ is the metric of spacetime and the indices μ and ν run from 0 to 3. Here, it is assumed that on the cosmic background, there exists a dominant stochastic magnetic field whose wavelengths are less than its curvature. Hence, the mean electromagnetic fields then become the source of Einstein equations [66]. The averaged electromagnetic fields are given by [2]:

$$\langle E \rangle = \langle B \rangle = 0, \langle E_i B_j \rangle = 0, \quad (4)$$

$$\langle E_i E_j \rangle = \frac{1}{3} E^2 g_{ij}, \langle B_i B_j \rangle = \frac{1}{3} B^2 g_{ij},$$

where $\langle \rangle$ denotes averaging brackets used for taking mean of volume and the indices i and j run from 1 to 3. The wavelength of radiation is considered to be lower than the volume and the volume smaller than the curvature.

However, the case of real nonlinear magnetic monopole is when $E^2 = 0$. Therefore, as obtained from equation (1), the energy density $\rho = -T_0^0$ and the pressure $p = T_i^i/3$ of the nonlinear monopole magnetic field is [5]

$$\rho_{\text{NED}} = -\mathcal{L}_{\text{NED}}, \quad (5)$$

$$p_{\text{NED}} = \mathcal{L}_{\text{NED}} - \frac{2B^2}{3} \frac{\partial \mathcal{L}_{\text{NED}}}{\partial \mathcal{F}}, \quad (6)$$

where the definition of \mathcal{L}_{NED} is given in Eq. (1) with $\mathcal{F} = B^2/2$.

From the above equations, we obtain the energy density equation ρ and pressure p as thus:

$$\begin{aligned} \rho &= \rho_{\text{NED}} = \frac{2^{-\alpha} (B^2)^{\alpha}}{4} \\ p &= p_{\text{NED}} = \frac{2^{-\alpha}}{12} (B^2)^{\alpha} (4\alpha - 3). \end{aligned} \quad (7)$$

3. Cosmology with variable G and nonlinear electrodynamics

In varying G theories, the action is still

$$S = \int d^4x \left(\sqrt{-g} \left(\frac{R}{16\pi G} + \mathcal{L}_{\text{NED}} \right) \right). \quad (8)$$

Taking the variation of the action with respect to the metric and ignoring surface terms leads to

$$G_{\mu\nu} - g_{\mu\nu} \Lambda = \frac{8\pi G}{c^4} T_{\mu\nu}. \quad (9)$$

In cosmological context, the Friedmann Robertson Walker metric is written as

$$ds^2 = -c^2 dt^2 + a(t)^2 \left[\frac{dr^2}{1 - Kr^2} + r^2 d\Omega \right], \quad (10)$$

where $a(t)$ is the scale factor, t is the comoving time and $K = 0, 1, -1$ represent a flat, closed and open FRW universe, respectively.

For the case of flat FRW ($K = 0$) and $c = 1$, the Einsteins equations are [27–29],

$$H^2 = \left(\frac{\dot{a}}{a} \right)^2 = \frac{8\pi G(t)}{3} \rho, \quad (11)$$

$$\frac{\ddot{a}}{a} = -\frac{4\pi G(t)}{3} (\rho + 3p). \quad (12)$$

where H represents the Hubble parameter and dot is the differentiation with respect to time.

However, the conservation equation that follows from (11)–(12) is for time variation in $G(t)$ is now [46, 47]:

$$\dot{\rho} + 3 \frac{\dot{a}}{a} (\rho + p) = -\rho \frac{\dot{G}}{G}, \quad (13)$$

where $\omega = \frac{p}{\rho}$ denotes the equation of state parameter for the dark energy.

Without diving into the dynamics of variable, we shall use Barrow's ansatz in which the gravitational constant G varies in form of the power-law of the scale factor as [46, 47]:

$$G = G_0 a^m, \quad (14)$$

where G_0 is a positive constant. Since we know that G increases with time, m must be positive. Furthermore, since G depends on the scale factor of the universe, its time derivatives \dot{G} must be greater than zero.

From the conservation equation (13), we obtain:

$$-\frac{\partial \mathcal{L}_{\text{NED}}}{2\partial \mathcal{F}} \cdot \left(\frac{d}{dt} ((B(t))^2) + 4 \frac{B(t)^2 \dot{a}}{a} \right) - \frac{\mathcal{L}_{\text{NED}} \dot{G}}{G} = 0. \quad (15)$$

The solution to the above equation gives an important relation between $B(t)$ and $a(t)$ as follows:

$$B(t) = a(t)^{-1/2 \frac{4\alpha+m}{\alpha}} B_0. \quad (16)$$

Conveniently, when written in terms of the scale factor, the evolution of energy density and pressure are given by:

$$\rho = \frac{2^{-\alpha}}{4} \left(B_0^2 a(t)^{-\frac{4\alpha+m}{\alpha}} \right)^{\alpha}, \quad (17)$$

$$p = \frac{2^{-\alpha}}{12} \left(B_0^2 a(t)^{-\frac{4\alpha+m}{\alpha}} \right)^{\alpha} (4\alpha - 3). \quad (18)$$

Then we have:

$$\rho + p = \frac{2^{-\alpha}}{3} \left(B_0^2 a(t)^{\frac{-4\alpha-m}{\alpha}} \right)^\alpha, \quad (19)$$

$$\rho + 3p = 2^{-1-\alpha} \left(B_0^2 a(t)^{\frac{-4\alpha-m}{\alpha}} \right)^\alpha (2\alpha - 1), \quad (20)$$

and the EoS parameter ω is

$$\omega = \frac{4}{3}\alpha - 1. \quad (21)$$

Equation (21) shows some interesting cases of the universe. It follows that at $\alpha = 0$, $\omega = -1$, is a cosmological constant case, at

$\alpha = 0.5$, $\omega = -1/3$ indicates case of dark energy and at $\alpha = 1$, $\omega = 1/3$ denotes the case of ultra-relativistic. The matter content of the universe is related to its acceleration equation:

$$\frac{\ddot{a}}{a} = -\frac{4\pi G(t)}{3}(\rho + 3p). \quad (22)$$

On checking the singularity in energy density and pressure at $a(t) \rightarrow 0$ and $a(t) \rightarrow \infty$, we obtain,

$$\lim_{a(t) \rightarrow 0} \rho(t) = \lim_{a(t) \rightarrow 0} p(t) = 0, \quad (23)$$

$$\lim_{a(t) \rightarrow \infty} \rho(t) = \lim_{a(t) \rightarrow \infty} p(t) = 0. \quad (24)$$

Using the Einstein's field equation and energy momentum tensor, the Ricci Scalar R which gives the curvature of spacetime is calculated as follows:

$$R = 8\pi G_0 a(t)^m (\rho - 3p). \quad (25)$$

The Ricci tensor squared $R_{\mu\nu}R^{\mu\nu}$ and the Kretschmann scalar $R_{\mu\nu\alpha\beta}R^{\mu\nu\alpha\beta}$ are also obtained [5] as:

$$R_{\mu\nu}R^{\mu\nu} = (8\pi G_0 a(t)^m)^2 (\rho^2 + 3p^2), \quad (26)$$

$$R_{\mu\nu\alpha\beta}R^{\mu\nu\alpha\beta} = (8\pi G_0 a(t)^m)^2 \left(\frac{5}{3}\rho^2 + 2\rho p + 3p^2 \right), \quad (27)$$

$$\lim_{a(t) \rightarrow 0} R(t) = \lim_{a(t) \rightarrow 0} R_{\mu\nu}R^{\mu\nu} = \lim_{a(t) \rightarrow 0} R_{\mu\nu\alpha\beta}R^{\mu\nu\alpha\beta} = 0. \quad (28)$$

The nature of the scale factor gives the behaviour of the curvature scalar. Taking the limit of the above equations as the universe accelerates at $a(t) \rightarrow 0$, we obtain no singularities in the curvature scalar, Ricci tensor and the Kretschmann scalar.

4. The evolution of the scale factor of the universe

The first Friedmann equation with variable $G(t)$ for the flat universe is given by

$$H^2 = \left(\frac{\dot{a}}{a} \right)^2 = \frac{8\pi G(t)}{3} \rho. \quad (29)$$

When a particle moves in one dimension in a potential $V(a)$, the equation of motion is

$$\dot{a}^2 + V(a) = 0. \quad (30)$$

The potential function

$$V(a) = -\frac{1}{3} 2^{1-\alpha} \pi G_0 (a(t))^{m+2} \left(B_0^2 (a(t))^{\frac{-4\alpha-m}{\alpha}} \right)^\alpha, \quad (31)$$

is negative and possesses a maximum at $a = a_c = -C_1$.

Using the (14) and (5), it becomes

$$\frac{-2^{1-\alpha} \pi G_0 a^{-4\alpha+2} B_0^{2\alpha} + 3\dot{a}^2}{3a^2} = 0, \quad (32)$$

then we find the scale factor $a(t)$ is equal to

$$a(t) \approx \frac{1}{2} \sqrt{B_0} 2^{\frac{3}{4\alpha}} 2^{\frac{3}{4\alpha}} \left(G_0 \alpha^2 (C_2 - t)^2 \right)^{\frac{1}{4\alpha}} 3^{-\frac{1}{4\alpha}} \pi^{\frac{1}{4\alpha}}. \quad (33)$$

Equation (33) shows that at $t \rightarrow 0$, the scale factor $a(t) \rightarrow$

$$\frac{1}{2} \sqrt{B_0} 2^{\frac{3}{4\alpha}} 2^{\frac{3}{4\alpha}} \left(G_0 \alpha^2 C_2^2 \right)^{\frac{1}{4\alpha}} 3^{-\frac{1}{4\alpha}} \pi^{\frac{1}{4\alpha}}. \quad (34)$$

This implies that the size of the universe $a(t)$ was never zero.

In the early universe, there is a de Sitter phase because of the nonlinear corrections to Maxwell's theory, as shown in Fig. 1. Thus, for $\alpha < 1$, the universe enters into its accelerating expansion phase which indicates the present dark energy dominated epoch. When $\alpha \geq 1$ presents a




Fig. 1 Plot of the scale factor a versus the time t (for $B_0 = G_0 = C_2 = m = 1$)

radiation-dominated era with the universe in the phase of decelerating expansion [68].

Introducing the quantity q (the deceleration parameter) [68], we described the expansion of the universe by:

$$q = -\frac{\ddot{a}a}{(\dot{a})^2} = 9/2 \frac{\rho + 3p}{a\rho}. \quad (35)$$

There is a inflation phase for $q < 0$ for $\alpha < 1$ and deceleration phase for $q > 0$ for $\alpha > 1$ shown in Fig. 2.

To estimate the amount of the inflation, we use the definition of e-foldings

$$N = \ln \frac{a(t_{end})}{a(t_{in})} \quad (36)$$

where t_{end} is the time inflation ends while t_{in} is the time it begins. For $N \simeq 70$ e-folding, the cosmic flatness and horizon problems can be resolved. Hence, we obtain the scale factor for beginning time of inflation (for $m = 1, \alpha = 1, G_0 = 1, B_0 = 1$):

$$a(t_{in}) = 3.46 \times 10^{-31}. \quad (37)$$

Using the second Friedmann equation Eq. (22) which is known as the acceleration equation for the universe:

$$\frac{\ddot{a}}{a} = \frac{(1 - 2\alpha)}{4\alpha^2 t^2}. \quad (38)$$

It is clear that acceleration stops at $\alpha = 0.5$.

5. Test of Causality

For any cosmological model to survive, it is an established that the speed of sound can not exceed the local speed of light, $c_s \leq 1$. The second requirement for stability is that the square of the speed of sound must be positive, i.e. $c_s^2 > 0$. In case the model is a classically stable one, [67] at $E = 0$, we obtain:

$$c_s^2 = \frac{dp}{d\rho} = \frac{dp/d\mathcal{F}}{d\rho/d\mathcal{F}} = -\frac{7}{3} + \frac{4}{3}\alpha. \quad (39)$$

A requirement of the classical stability $c_s^2 > 0$ is $\alpha > \frac{7}{4}$, and the causality $c_s \leq 1$ is $\alpha \leq \frac{5}{2}$. Hence, the best value of α for both stability conditions is $\frac{5}{2} \geq \alpha > \frac{7}{4}$.

6. Conclusion

This work studied cosmology with a varying gravitational constant G and NED in a flat FRW universe. Under changes to the scale factor, the evolution of the magnetic field reduced to $B(t) = a(t)^{-1/2} B_0$, as obtained in [12] where $m = 0$. The evolution of the scale factor shows that the models give an accelerating, expanding universe with

$$a(t) \approx \frac{1}{2} \sqrt{B_0} 2^{\frac{3}{2\alpha}} 2^{\frac{3}{4\alpha}} \left(G_0 \alpha^2 (C_2 - t)^2 \right)^{\frac{1}{4\alpha}} 3^{-\frac{1}{4\alpha}} \pi^{\frac{1}{4\alpha}}, \quad (40)$$

where B_0 represents the magnetic induction field at present time t_0 , and α is a free parameter presented in Fig. 1. The value of the scale factor obtained indicates that the size of the universe was never zero. By incorporating NED and variable gravitational constant, we present a model free from the initial big bang singularity and inflation. As observed in equations (23), (24), and (28), the models portray no singularity in the energy density, pressure, and curvature terms, respectively. For $\alpha \geq 1$, the universe is in a radiation-dominated era with the phase of decelerating expansion. However, when $\alpha = 0.5$, $\omega = -1/3$ indicates a transition into the present epoch of dark energy with accelerating expansion. Furthermore, we studied the stability of this model and observed that it depends of the constant α and is classically stable for $\frac{5}{2} \geq \alpha > \frac{7}{4}$. In future studies, we intend to investigate the evolution of the universe with both varying G and c in NED.




Fig. 2 Plot of the deceleration parameter q versus the scale factor a (for $B_0 = G_0 = C_2 = m = 1$)

References

- [1] S Capozziello and V Faraoni *Beyond Einstein Gravity: A Survey of Gravitational Theories for Cosmology and Astrophysics Springer Science+Business Media B V* New York, 2011).
- [2] V A De Lorenci, R Klippert, M Novello and J M Salim *Phys. Rev. D* **65** 063501 (2002).
- [3] D N Vollick *Phys. Rev. D* **78** 063524 (2008).
- [4] S I Kruglov *Phys. Rev. D* **92** 123523 (2015).
- [5] A Övgün *Eur. Phys. J. C* **77** 105 (2017).
- [6] A Övgün, G Leon, J Magana and K Jusufi *Eur. Phys. J. C* **78** 462 (2018).
- [7] G Otalora, A Övgün, J Saavedra and N Videla *JCAP* **06** 003 (2018).
- [8] P Sarkar, P K Das and G C Samanta [arXiv:2005.05568 [hep-ph]].
- [9] A Golovnev, V Mukhanov and V Vanchurin *JCAP* **0806** 009 (2008).
- [10] M Novello, et al. *Phys.Rev. D* **69** 127301 (2004).
- [11] M Novello, E Goulart, J M Salim and S E Perez Bergliaffa *Class. Quant. Grav.* **24** 3021 (2007).
- [12] S I Kruglov *Int. J. Mod. Phys. A* **32** 1750071 (2017).
- [13] S I Kruglov *Annals Phys.* **353** 299 (2014).
- [14] S I Kruglov *Int. J. Mod. Phys. D* **25** 1640002 (2016).
- [15] S I Kruglov *Int. J. Mod. Phys. A* **31** 1650058 (2016).
- [16] S Kruglov *Eur. Phys. J. Plus* **135** 370 (2020).
- [17] M Sharif and S Mumtaz *Eur. Phys. J. C* **77** 136 (2017).
- [18] L Campanelli, P Cea, G L Fogli and L Tedesco *Phys. Rev. D* **77** 043001 (2008).
- [19] R Garcia-Salcedo et al. *Int. J. Mod. Phys. A* **15** 4341 (2000).
- [20] N J Poplawski *Gen. Rel. Grav.* **44**, 1007 (2012).
- [21] N Poplawski *Astrophys. J.* **832** 96 (2016).
- [22] A Öztaş, E Dil and M Smith *Mon. Not. Roy. Astron. Soc.* **476** 451 (2018).
- [23] E Dil, A M Oztas and E Dil *Astron. Comput.* **28** 100295 (2019).
- [24] F V Mukhanov et al. *Phys. Rev. Lett.* **68** 1969 (1992).
- [25] E N Saridakis and M Tsoukalas *Phys. Rev. D* **93** 124032 (2016).
- [26] H Sheikahmadi, E N Saridakis, A Aghamohammadi and K Saaidi *JCAP* **10** 021 (2016).
- [27] C S Camara et al. *Phys.Rev. D* **69** 123504 (2004).
- [28] R Durrer and A Neronov *Astron Astrophys Rev* **21** 62 (2013).
- [29] K E Kunze *Plasma Phys. Control. Fusion* **55** 124026 (2013).
- [30] H Azri and D Demir *Phys. Rev. D* **95** 124007 (2017).
- [31] H Azri and D Demir *Phys. Rev. D* **97** 044025 (2018).
- [32] H Azri and A Bounames *Int. J. Mod. Phys. D* **26** 1750060 (2017).
- [33] A Hernandez-Almada, M A Garcia-Aspeitia, J Magana and V Motta *Phys. Rev. D* **101** 063516 (2020).
- [34] M A Garcia-Aspeitia, C Martinez-Robles, A Hernandez-Almada, J Magana and V Motta *Phys. Rev. D* **99** 123525 (2019).
- [35] J A Vazquez, L E Padilla and T Matos *Revista Mexicana de Física E* 2020 [arXiv:1810.09934 [astro-ph.CO]].
- [36] O Sert *Mod. Phys. Lett. A* **35** 2050037 (2019).
- [37] M Adak, O Akarsu, T Dereli and O Sert *JCAP* **11** 026 (2017).
- [38] O Okcu, C Corda and E Aydiner *EPL* **129** 50002 (2020).
- [39] M Cruz, F Izaurieta and S Lepe *Eur. Phys. J. C* **80** 559 (2020).
- [40] M Cruz and S Lepe *Class. Quant. Grav.* **35** 155013 (2018).
- [41] M Cruz, N Cruz and S Lepe *Phys. Rev. D* **96** 124020 (2017).
- [42] R Cordero, M Cruz, A Molgado and E Rojas *Gen. Rel. Grav.* **46** 1761 (2014).
- [43] E Elizalde et al. *Phys. Lett. B* **574** 1 (2003).
- [44] C Quercellini, M Bruni, A Balbi, D Pietrobon *Phys. Rev. D* **78** 063527 (2008).
- [45] M Born and L Infeld *Proc. Roy. Soc. Lond. A* **144** 425 (1934).
- [46] J D Barrow and J Magueijo *Phys. Lett. B* **443** 104 (1998).
- [47] A Albrecht and J Magueijo *Phys. Rev. D* **59** 043516 (1999).
- [48] A G Cohen, D B Kaplan and A E Nelson *Phys. Rev. Lett.* **82** 4971 (1999).
- [49] J Moffat *Eur. Phys. J. C* **76** 130 (2016).
- [50] K Leszczynska *Universe* **3** 46 (2017).
- [51] C Camara, R Nascimento, J Carvalho and M De Garcia Maia *Int. J. Mod. Phys. D* **16** 433 (2007).
- [52] L Kantha *Adv. Astron.* **2016** 9743970 (2016).
- [53] S Nesseris L Perivolaropoulos *Phys.Rev. D* **73** 103511 (2006).
- [54] H Weyl *Naturwissenschaften* **22** 145 (1934).
- [55] P A M Dirac *Proc. R. Soc. A* **165** 199 (1938).
- [56] A K Singha and U Debnath *Int. J. Mod. Phys. D* **16** 117 (2007).
- [57] M S Singh and S S Singh *Turk J Phys* **42** 198 (2018).
- [58] R Dungan and H B Prosper [arXiv:0909.5416 [astro-ph.CO]].
- [59] M Malekjani, J Lu, N Nazari-Pooya, L Xu, D Mohammad-Zadeh Jassur and M Honari-Jafarpour *Astrophys. Space Sci.* **360** 24 (2015).
- [60] A Sheykhi *Phys. Scripta* **85** 045901 (2012).
- [61] J Lu, L Xu, H Tan and S Gao *Phys. Rev. D* **89** 063526 (2014).
- [62] M Sharif and A Jawad *Eur. Phys. J. C* **72** 1901 (2012).
- [63] J Lu, E N Saridakis, M R Setare and L Xu *JCAP* **1003**, 031 (2010).
- [64] A Sheykhi and M R Setare *Int. J. Theor. Phys.* **49** 2777 (2010).
- [65] B Borah and M Ansari *Int. J. Theor. Phys.* **53** 1217 (2014).
- [66] R Tolman, P Ehrenfest *Phys.Rev.* **36** 1791 (1930).
- [67] R Garcia-Salcedo, T Gonzalez, I Quiros *Phys.Rev. D* **89** 084047 (2014).
- [68] S M Carroll [arXiv:gr-qc/9712019 [gr-qc]].

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.